

## Optical approaches to examining the dynamics of dissolved organic carbon in optically complex inland waters

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2012 Environ. Res. Lett. 7 034014

(<http://iopscience.iop.org/1748-9326/7/3/034014>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 159.226.73.206

The article was downloaded on 09/07/2013 at 02:28

Please note that [terms and conditions apply](#).

# Optical approaches to examining the dynamics of dissolved organic carbon in optically complex inland waters

Guangjia Jiang<sup>1,2</sup>, Ronghua Ma<sup>1,4</sup>, Steven A Loiselle<sup>3</sup> and Hongtao Duan<sup>1</sup>

<sup>1</sup> State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 210008, People's Republic of China

<sup>2</sup> Graduate University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

<sup>3</sup> Dipartimento Farmaco Chimico Tecnologico, CSGI, University of Siena, Siena, Italy

E-mail: [gjiang2011@gmail.com](mailto:gjiang2011@gmail.com), [mrhua2002@niglas.ac.cn](mailto:mrhua2002@niglas.ac.cn), [steven.loiselle@unisi.it](mailto:steven.loiselle@unisi.it) and [htduan@niglas.ac.cn](mailto:htduan@niglas.ac.cn)

Received 15 May 2012

Accepted for publication 7 August 2012

Published 16 August 2012

Online at [stacks.iop.org/ERL/7/034014](http://stacks.iop.org/ERL/7/034014)

## Abstract

Optical approaches were developed to examine the relationship between the optically active and the optically inactive fractions of dissolved organic carbon in inland waters. A multiple linear regression model was developed on the basis of an extensive dataset from Taihu Lake, and validated employing data from another large shallow eutrophic lake (Chaohu Lake) in southern China. The model was used to estimate the concentration of dissolved organic matter ( $n = 191$ ) using the absorption characteristics of its chromophoric fraction with a high correlation coefficient ( $R^2 = 0.62$ ) and a low root mean squared error (RMSE = 9.67%). This intra-ecosystem validity allows us to improve our understanding of carbon dynamics using optical remote sensing approaches for these optically complex lakes, where multiple sources and sinks of dissolved organic matter were present.

**Keywords:** CDOM, DOC, lake, optical properties, remote sensing, case 2 waters

## 1. Introduction

In recent decades, increasing attention has been paid to the dynamics of organic carbon in natural waters. As dissolved organic carbon (DOC) is the largest pool of reduced organic carbon in most aquatic ecosystems (Hedges 2002), an understanding of its spatial and temporal dynamics is fundamental to monitor carbon cycling in marine and lake environments (Chen *et al* 2004). DOC is a major energy source for microbially based aquatic food webs


(Deegan and Garritt 1997) and plays an important role in the biogeochemical dynamics and optical conditions of many lakes and rivers. Spectral attenuation by chromophoric dissolved organic matter (CDOM), the colored fraction of DOC, can modify primary productivity as well as reduce UV related stress on aquatic biota (Nieke *et al* 1997).

Identifying a relationship between the chromophoric and non-chromophoric portions of dissolved organic matter allows for the optical analysis of aquatic carbon cycles, particularly important in large lakes and marine environments. This relationship has been investigated in rivers (Del Castillo and Miller 2008), estuarial waters (Spencer *et al* 2009) and coastal systems (Fichot and Benner 2011). Many studies indicate that the spectral absorption characteristics of CDOM ( $a_g(\lambda)$ ) and the concentration of DOC ( $C_{DOC}$ ) vary independently (Vodacek *et al* 1997, Loiselle *et al* 2009b). However,



Content from this work may be used under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 licence](http://creativecommons.org/licenses/by-nc-sa/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.


<sup>4</sup> Author to whom any correspondence should be addressed.



**Figure 1.** Geographical location of sampling sites in Taihu Lake and Chaohu Lake. For Taihu Lake, overlaid points in Meiliang Bay and Zhushan Bay result from sampling in the same positions in different cruises.

recent studies in coastal waters have shown that where a single DOC source is dominant, DOC concentrations can be estimated using a log-linearized ecosystem specific relationship using two CDOM absorption wavelengths,  $a_g(275)$  and  $a_g(295)$  (Fichot and Benner 2011). Inland waters, however, usually have multiple processes that modify the relation between the optical and chemical properties of DOC (dilution, photodegradation, flocculation) (Chen *et al* 2004, Loiselle *et al* 2010).

Changes in CDOM absorption characteristics derive directly from solar exposure and indirectly oxidation of the aromatic compounds (Reche *et al* 2000, Loiselle *et al* 2012), which cause losses in humification and aromaticity (Weishaar *et al* 2003). The molecular implications of photobleaching differ depending on the spectral characteristics of the incident solar irradiance. In the ultraviolet B wavelengths (UVB, 280–315 nm), absorption losses of CDOM are related to the degradation of the olefinic components (Loiselle *et al* 2009b). Exposure to UVA (315–400 nm) has been associated to modifications in the CDOM aromatic fraction (Corin *et al* 1996, Ma and Green 2004). Changes in absorption in the visible wavelengths have been associated with a



**Figure 2.** Spectral slopes against wavelength from 250 to 400 nm calculated using non-linear regression at a 20 nm interval in the four cruises in Taihu Lake.

disruption in the intramolecular charge transfer that is related to the featureless absorption spectrum typical of natural CDOM (Osburn *et al* 2009).

Changes in the optical properties of CDOM can be characterized using individual absorption wavelengths, specific UV absorption  $a_g^*(\lambda) = a_g(\lambda)/C_{DOC}$  (Weishaar *et al* 2003), the ratio of absorptions  $a_R = a_g(\lambda_1)/a_g(\lambda_2)$  (Helms *et al* 2008) and the absorption spectral slope ( $S$ ,  $\text{nm}^{-1}$ ):  $a_g(\lambda) = a_g(\lambda_0) \exp[S(\lambda_0 - \lambda)]$  where  $a_g(\lambda)$  is the CDOM absorption at wavelength  $\lambda$  and  $a_g(\lambda_0)$  is the CDOM absorption at a reference wavelength  $\lambda_0$  (Bricaud *et al* 1981). Changes in  $S$  have been shown to reflect changes in the molecular structure and average molecular weight of DOM (Carder *et al* 1989).

Through improving our understanding of the linkages between  $C_{DOC}$  and its spectral properties, new insights into organic carbon dynamics can be gained, including the photochemical processes that modify the structure, reactivity and bio-lability of DOC in aquatic ecosystems. In the present study, we explore a modeling approach to estimate DOC from CDOM absorption using an extensive bio-optic dataset. This model is then applied in two shallow inland lakes.

## 2. Material and methods

Data from two shallow lakes (figure 1) were used in this study. Taihu Lake, the third largest freshwater lake in China with an average depth of 1.9 m, is characterized by severe summertime cyanobacteria blooms (Duan *et al* 2009). Chaohu Lake is another shallow and large hypereutrophic lake in south China, with similar eutrophication problems (Xie *et al* 2005).

Water samples (192) were collected from Taihu Lake at 5 cm depth in April 2010, March, May and August 2011 (figure 1 and table 1). Discrete samples were immediately filtered through precombusted (6 h at 450 °C) 47 mm Whatman GF/F glass fiber filters (0.70  $\mu\text{m}$ ) at a low




Figure 3. Relationships between slopes (275–295 nm, 295–365 nm, 275–365 nm) and  $a_R$  and  $a_g^*(365)$ .




Figure 4. Relationship between CDOM absorptions and DOC concentration in Taihu Lake: (a) 201 004; (b) 201 103; (c) 201 105; (d) 201 108.

Table 1. Descriptive statistics of averages of  $a_g(365)$ ,  $C_{DOC}$ ,  $a_g^*(365)$  and  $a_R$  from measurement campaigns in Taihu Lake and Chaohu Lake.

Sampling date	Proxy	Study area	Sampling points	$a_g(365)$ ( $m^{-1}$ )	$C_{DOC}$ ( $mg\ l^{-1}$ )	$a_g^*(365)$ ( $m^2\ g^{-1}$ )	$a_R$
23 April–2 May 2010	201 004	Taihu Lake	84	$2.30 \pm 0.83$	$4.77 \pm 0.62$	$0.47 \pm 0.12$	$8.79 \pm 1.34$
24–25 March 2011	201 103	Meiliang Bay Zhushan Bay	26	$3.27 \pm 1.46$	$4.79 \pm 0.80$	$0.67 \pm 0.25$	$7.99 \pm 1.67$
1–8 May 2011	201 105	Taihu Lake	54	$3.31 \pm 1.82$	$3.70 \pm 0.87$	$0.91 \pm 0.45$	$7.16 \pm 2.09$
31 August–2 September 2011	201 108	Meiliang Bay Zhushan Bay	27	$3.26 \pm 0.75$	$4.75 \pm 0.51$	$0.68 \pm 0.14$	$6.86 \pm 0.97$
15–16 October 2009	200 910	Chaohu Lake	35	$5.92 \pm 6.67$	$9.50 \pm 12.88$	$0.64 \pm 0.37$	$9.46 \pm 1.23$

vacuum for DOC analysis ( $C_{DOC}$ ), using a Shimadzu TOC-5000A analyzer following the protocol of Chen *et al* (2004). An aliquot of each sample was filtered through a  $0.22\ \mu m$  Whatman Nuclepore filter pretreated by 10% dilute hydrochloric acid. The filtrate was stored in the dark at a low temperature ( $4-8\ ^\circ C$ ) until analysis. The CDOM absorbance spectra were measured on a UV-2401 spectrophotometer with 1 cm quartz cuvettes, after the filtrate reached the room temperature ( $20\ ^\circ C$ ). Milli-Q water was used as a blank and reference. Absorbance ( $A(\lambda)$ ) from 200 to 800 nm was

determined for each sample and converted to the absorption coefficients as  $a_g(\lambda) = 2.303A(\lambda)/l$ , where  $l$  was the cuvette path length. Baseline correction was performed on the basis of the average absorption coefficients from 700 to 800 nm according to Green and Blough (1994).

Specific UV absorption coefficients,  $a_g^*(\lambda)(m^2\ g^{-1})$ , were calculated by dividing the CDOM absorption at  $\lambda = 365\ nm$  by  $C_{DOC}$ . This wavelength was chosen to be consistent with other studies which examine both CDOM absorption and fluorescence (Bari and Farooq 1985). Absorption ratios,

$a_R$ , were determined using CDOM absorptions at 255 and 365 nm:  $a_R = a_g(255):a_g(365)$  as given by Helms *et al* (2008).

The distribution of absorption spectral slope was calculated using a non-linear fitting method based on the least squares regression analysis from 250 to 400 nm at a 20 nm wavelength interval following Loiselle *et al* (2009b).

A model validation dataset was constructed using DOC and CDOM measurements on samples obtained in Chaohu Lake (30 stations) in October 2009 (table 1). Sampling and analysis were made following the same methods in both lakes.

### 3. Results and discussion

#### 3.1. Characteristics of DOC and CDOM in Taihu and Chaohu Lakes


In Taihu Lake, CDOM absorption at 365 nm averaged  $2.3 \text{ m}^{-1}$  in 2010 and  $3.3 \text{ m}^{-1}$  in 2011, while DOC concentrations ranged from 3.8 to  $4.8 \text{ mg l}^{-1}$  (table 1). The two northern bays, Meiling and Zhushan, often linked to algal blooms (Duan *et al* 2009), had similar absorption values to that of whole lake in 2011, but with much higher  $C_{\text{DOC}}$  values. The Chaohu Lake dataset presented higher  $a_g(365)$  and  $C_{\text{DOC}}$ . The carbon-specific UV absorption,  $a_g^*(365)$ , was highest in the Taihu Lake May 2011 dataset, averaging almost twice that of the Taihu Lake 2010 dataset obtained in the same period. The two northern bay datasets, obtained in March and September 2011 showed similar values. The average absorption ratio value ( $a_R$ ) was highest in the Chaohu Lake and 2010 Taihu Lake datasets.

The distribution of the absorption spectral slopes showed the spectral variability of  $S_\lambda$  and significant differences between sampling campaigns (figure 2). A significant and negative relationship was found between the absorption spectral slope ( $S_{275-295}$ ,  $S_{295-365}$ ,  $S_{275-365}$ ) and the carbon-specific UV absorption coefficients,  $a_g^*(365)$  (figure 3). As expected, a strong and positive correlation was observed between the absorption spectral slope intervals and  $a_R$ . Both terms have been used as indicators of intramolecular structure and weight of DOM (Helms *et al* 2008). Furthermore, the absorption spectral slope is clearly a logarithmic form of the absorption ratio. In the present dataset, these parameters,  $a_R$  (or  $S$ ) and  $a_g^*(365)$  were shown to provide comparable information on the CDOM present.

#### 3.2. Estimation of DOC from CDOM


The direct correlation between  $C_{\text{DOC}}$  and CDOM absorption was spectrally variable and generally poor for most datasets. The highest correlation was found to occur at lower wavelengths (figure 4).

The relationship between multiple wavelengths was explored using a multiple regression approach. In the recent work by Fichot and Benner (2011), a multiple regression analysis of the logarithm of CDOM absorption at 275 and 295 nm was used to estimate DOC concentrations in a coastal waters ecosystem. A similar approach, using the extensive



**Figure 5.** Determination of DOC concentrations from CDOM absorptions using multiple linear regressions of DOC concentrations against two and three CDOM absorptions.

datasets from Taihu Lake was used to determine dataset specific and lake specific algorithms following:  $C_{\text{DOC}} = \alpha + \beta a_g(275) + \gamma a_g(295)$ , where  $\alpha$ ,  $\beta$  and  $\gamma$  are regression coefficients. The resulting algorithms (figure 5) provided good results for each dataset ( $0.71 < R^2 < 0.85$ ), but did not present a strong correlation when the Taihu datasets were combined ( $R^2 = 0.49$ ,  $\text{RMSE} = 13.57\%$ ).



**Figure 6.** Validation of the Taihu Lake two wavelength model (a) and three wavelength model (b) to estimate DOC concentrations in Chaohu Lake.

We then examined a three wavelengths regression using CDOM absorption at 275, 295 and 365 nm:

$$C_{DOC} = \delta + \varepsilon a_g(275) + \zeta a_g(295) + \eta a_g(365) \quad (1)$$

where  $\delta$ ,  $\varepsilon$ ,  $\zeta$  and  $\eta$  are regression coefficients. Longer UV wavelengths were chosen to provide information related to changes in structure and sources of organic matter. The selection of these wavelengths was also related to absorption wavelengths used in other studies. The three wavelength algorithms provided significantly higher correlations between the estimated and measured DOC for all but one measurement campaign (201105) (figure 5). Using the complete Taihu Lake dataset, the resulting three wavelength algorithm showed a significantly higher correlation with DOC ( $R^2 = 0.62$ , RMSE = 9.67%).

$$C_{DOC} = 1.81 + 0.72a_g(275) - 0.79a_g(295) \quad (2)$$


$$C_{DOC} = 1.48 - 0.03a_g(275) + 0.60a_g(295) - 0.99a_g(365). \quad (3)$$

To examine the specificity of these algorithms developed for Taihu Lake, we tested the two regressions employing the Chaohu Lake dataset. The two wavelength model failed to show a significant estimation of  $C_{DOC}$  in Chaohu Lake. However, the three waveband model, optimized for Lake Taihu, provided a good estimate of Chaohu Lake  $C_{DOC}$  (figure 6). The RMS error was 15.59% and the errors were uniformly distributed around the 1:1 line. It should be noted that many samples are clustered around 4.5 mg l<sup>-1</sup>, but the relationship remains robust when part of these values are removed.

A separate three band model was developed using the dataset in Chaohu Lake, yielding the regression coefficients  $\delta = 2.02$ ,  $\varepsilon = -0.07$ ,  $\zeta = 0.49$  and  $\eta = -0.49$ , which were very similar to those of the Taihu Lake dataset ( $R^2 = 0.75$ , RMSE = 15.05%) (figure 7).

#### 4. Conclusion

The possibility of a direct relationship between DOC concentration and CDOM absorption is limited to ecosystems



**Figure 7.** DOC concentration estimated using the three wavelength model based on Chaohu Lake data.

where a single DOC source undergoes dilution or conservative mixing (Del Vecchio and Blough 2004). The lack of such a correlation indicates that multiple sources and sinks are present. Lake Taihu, similar to most inland water bodies, presents a combination of DOC sources and sinks, which vary spatially as well as temporally. In most productive inland ecosystems, both terrestrial and algal-derived DOC sources are present. In shallow unstratified lakes, upwelling and resuspension also bring DOC into the surface waters. Bacterial activity and photodegradation will usually dominate loss processes in large lake ecosystems. Both mechanisms have spatial and temporal variability in Taihu Lake, which has been confirmed in the spatial and temporal variability in the  $S_\lambda$  (Loiselle *et al* 2009a) and  $a_g^*(365)$ .

Based on the temporally and spatially distinct datasets from Taihu Lake, a multiple linear regression approach using three absorption wavelengths provided a viable estimation approach to estimate DOC concentrations, in particular when a heterogeneous dataset was considered. Furthermore, the same model was found to provide a good estimate of DOC concentrations in a different shallow lake ecosystem. The results show a high determination coefficient  $R^2 = 0.73$  and

a low RMS error 15.59%, indicating that the model was significantly robust for exploring DOC dynamics in this second lake.

Although the methodology may be applied to other aquatic ecosystems, re-parameterization of the model may be required where differences in geochemical and biological processes influence the optical properties of DOM. The links between the optical and chemical properties of DOM improve our understanding of the temporal and spatial variability of its molecular characteristics. Furthermore, such information makes it possible to examine relationships between optically active and optically inactive components of the DOM and to determine spectrally distinct degradation rates in relation to exposure to solar radiation (photodegradation), a major transformation process of dissolved organic carbon.

## Acknowledgments

We gratefully acknowledge the financial support of the Knowledge Innovation Program of the Chinese Academy of Sciences (Grant Nos KZCX2-YW-QN311, KZCX2-EW-QN308) and National Natural Science Foundation of China (Grant Nos 40871168, 40801137). We thank Wen Su, Linlin Shang, Lin Zhou, Chunguang Lv, Jiawang Rao and Chenlu Zhao for participating in the field campaigns. We are also indebted to the two anonymous reviewers for their comments which greatly improved this manuscript.

## References

- Bari A and Farooq S 1985 Measurement of wastewater treatment efficiency by fluorescence and UV absorbance *Environ. Monit. Assess.* **5** 423–34
- Bricaud A, Morel A and Prieur L 1981 Absorption by dissolved organic matter of the sea (yellow substance) in the UV and visible domains *Limnol. Oceanogr.* **26** 43–53
- Carder K L, Steward R G, Harvey G R and Ortner P B 1989 Marine humic and fulvic acids: their effects on remote sensing of ocean chlorophyll *Limnol. Oceanogr.* **34** 68–81
- Chen Z, Li Y and Pan J 2004 Distributions of colored dissolved organic matter and dissolved organic carbon in the Pearl River Estuary, China *Cont. Shelf Res.* **24** 1845–56
- Corin N, Backlund P and Kulovaara M 1996 Degradation products formed during UV-irradiation of humic waters *Chemosphere* **33** 245–55
- Deegan L A and Garritt R H 1997 Evidence for spatial variability in estuarine food webs *Mar. Ecol. Prog. Ser.* **147** 31–47
- Del Castillo C E and Miller R L 2008 On the use of ocean color remote sensing to measure the transport of dissolved organic carbon by the Mississippi River Plume *Remote Sens. Environ.* **112** 836–44
- Del Vecchio R and Blough N V 2004 Spatial and seasonal distribution of chromophoric dissolved organic matter and dissolved organic carbon in the Middle Atlantic Bight *Mar. Chem.* **89** 169–87
- Duan H T, Ma R H, Xu X F, Kong F X, Zhang S X, Kong W J, Hao J Y and Shang L L 2009 Two-decade reconstruction of algal blooms in China's Lake Taihu *Environ. Sci. Technol.* **43** 3522–8
- Fichot C G and Benner R 2011 A novel method to estimate DOC concentrations from CDOM absorption coefficients in coastal waters *Geophys. Res. Lett.* **38** L03610
- Green S A and Blough N V 1994 Optical absorption and fluorescence properties of chromophoric dissolved organic matter in natural waters *Limnol. Oceanogr.* **39** 1903–16
- Hedges J I 2002 Why dissolved organics matter *Biogeochemistry of Marine Dissolved Organic Matter* ed D A Hansell and C A Carlson (San Diego, CA: Academic) pp 1–33
- Helms J R, Stubbins A, Ritchie J D, Minor E C, Kieber D J and Mopper K 2008 Absorption spectral slopes and slope ratios as indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter *Limnol. Oceanogr.* **53** 955–69
- Loiselle S A, Azza N, Gichuki J, Bracchini L, Tognazzi A, Dattilo A M, Rossi C and Cozar A 2010 Spatial dynamics of chromophoric dissolved organic matter in nearshore waters of Lake Victoria *Aquat. Ecosyst. Health Manag.* **13** 185–95
- Loiselle S A, Bracchini L, C  zar A, Dattilo A M, Tognazzi A and Rossi C 2009a Variability in photobleaching rates and their related impacts on optical conditions in subtropical lakes *J. Photochem. Photobiol. B* **95** 129–37
- Loiselle S A, Bracchini L, Dattilo A M, Ricci M, Tognazzi A and C  zar A 2009b Optical characterization of chromophoric dissolved organic matter using wavelength distribution of absorption spectral slopes *Limnol. Oceanogr.* **54** 590–7
- Loiselle S A, Vione D, Minero C, Maurino V, Tognazzi A, Dattilo A, Rossi C and Bracchini L 2012 Chemical and optical phototransformation of dissolved organic matter *Water Res.* **46** 3197–207
- Ma X D and Green S A 2004 Photochemical transformation of dissolved organic carbon in lake superior—an *in situ* experiment *J. Great Lakes Res.* **30** 97–112
- Nieke B, Reuter R, Heuermann R, Wang H, Babin M and Theriault J C 1997 Light absorption and fluorescence properties of chromophoric dissolved organic matter (CDOM), in the St. Lawrence Estuary (Case 2 waters) *Cont. Shelf Res.* **17** 235–52
- Osburn C L, O'Sullivan D W and Boyd T J 2009 Increases in the longwave photobleaching of chromophoric dissolved organic matter in coastal waters *Limnol. Oceanogr.* **54** 145–59
- Reche I, Pace M L and Cole J J 2000 Modeled effects of dissolved organic carbon and solar spectra on photobleaching in lake ecosystems *Ecosystems* **3** 419–32
- Spencer R G M, Aiken G R, Butler K D, Domblaser M M, Striegl R G and Hernes P J 2009 Utilizing chromophoric dissolved organic matter measurements to derive export and reactivity of dissolved organic carbon exported to the Arctic Ocean: a case study of the Yukon River, Alaska *Geophys. Res. Lett.* **36** L06401
- Vodacek A, Blough N V, DeGrandpre M D, Peltzer E T and Nelson R K 1997 Seasonal variation of CDOM and DOC in the Middle Atlantic Bight: terrestrial inputs and photooxidation *Limnol. Oceanogr.* **42** 674–86
- Weishaar J L, Aiken G R, Bergamaschi B A, Fram M S, Fugii R and Mopper K 2003 Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon *Environ. Sci. Technol.* **37** 4702–8
- Xie L Q, Xie P, Guo L G, Li L, Miyabara Y and Park H 2005 Organ distribution and bioaccumulation of microcystins in freshwater fish at different trophic levels from the eutrophic Lake Chaohu, China *Environ. Toxicol.* **20** 293–300